

TABLE 1 COMPARISON OF *SMINTHURINUS* FROM CENTRAL AND SOUTH AMERICA

SPECIES	LENGTH mm	MUCRO EDGES SERRATE	PAPILLA ANT III (Ant. Ratio)	TENENT HAIRS I, II, III	UNGUAL PSEUDON YCHIA	TUNIC A	DENS		TENACULAR		UNGUIC ULAR THEETH	SUBANAL APPEND PALMATE	Bulb
							TOTAL SETAE E4 /	SETAE	RAMI /SETAE	SETAE			
antennalis	0.31	0	1 1:1.1;1.4;3	0,0,0	%	?	?	13 (15 ?)	3	0	+	?	+
australis	2.4	0	1 1:1.8;2.5;6.9	3,3,3	+ ?	- ?		23	3	1	-	?	-
bullai	1.0	0	1 ? 1:1.7;2.1;5.7	2,3,3	?	?		?	3	0	?	?	?
castagninoi	0.71-0.82	2	1	3,3,3	?	?		?	3	1	?	+	?
inexcussus	0.57-0.63	1	1 1:1.7;2.4;6	3,3,3	+?	+	-	13	3	1	-	+	-
insularis	0.81	1	1? 1:1.9;2.6;6.2	3,3,3	?	?	-	13	3	2	?	?	?
molinai	1.0-1.2	0	1 1:2;2.8;6.6	4-5,4-5,4-5	+	?	?	19	ι	2	?	-	?
nunezi	1.0	2	1 *1:1.5:2:5	3,3,2	+	+	+	18	ι	1	?	+	?
operosus	1.3	1	1 1:2;2.8;7.5	5,5,5	-	-	?	16	3	2	+	+	+
packardi	0.70	0	1 1:2;2.8;6.6	4,4,4	?	?	?	19	3	2	-	+	-
patagonicus	0.80	2	1 1:1.7;2;3.6	5,5,5	?	-	-	15	3	1	+	+	+
ayi	0.3-1.2	2	1-2 1:2:1.7:2.5?	5,5,5	+	-	-	23	3	1-2	+	+	-
tuberculatus	?	2 week	1	0,0,0	-	-	-	18*	3	?	?	+	-
tucamanensis	?	2 strong	4	4,4,3	-	-	-	20	3	?	?	+	-
woringeri	?	1	1	2,3,3	+	+	?	18	3	?	?	+	-
harryius	0.90	2	4	7,6,6	+	+	?	17	3	2	-	+	-

* Four spines on each side.